

Marking Scheme
Strictly Confidential
(For Internal and Restricted use only)
Secondary School Examination, 2023
SUBJECT NAME: MATHEMATICS (BASIC)
(PAPER CODE 430/5/3)

General Instructions: -

1	You are aware that evaluation is the most important process in the actual and correct assessment of the candidates. A small mistake in evaluation may lead to serious problems which may affect the future of the candidates, education system and teaching profession. To avoid mistakes, it is requested that before starting evaluation, you must read and understand the spot evaluation guidelines carefully.
2	“Evaluation policy is a confidential policy as it is related to the confidentiality of the examinations conducted, Evaluation done and several other aspects. Its’ leakage to public in any manner could lead to derailment of the examination system and affect the life and future of millions of candidates. Sharing this policy/document to anyone, publishing in any magazine and printing in News Paper/Website etc may invite action under various rules of the Board and IPC.”
3	Evaluation is to be done as per instructions provided in the Marking Scheme. It should not be done according to one’s own interpretation or any other consideration. Marking Scheme should be strictly adhered to and religiously followed. However, while evaluating, answers which are based on latest information or knowledge and/or are innovative, they may be assessed for their correctness otherwise and due marks be awarded to them. In class-X, while evaluating two competency-based questions, please try to understand given answer and even if reply is not from marking scheme but correct competency is enumerated by the candidate, due marks should be awarded.
4	The Marking scheme carries only suggested value points for the answers. These are in the nature of Guidelines only and do not constitute the complete answer. The students can have their own expression and if the expression is correct, the due marks should be awarded accordingly.
5	The Head-Examiner must go through the first five answer books evaluated by each evaluator on the first day, to ensure that evaluation has been carried out as per the instructions given in the Marking Scheme. If there is any variation, the same should be zero after deliberation and discussion. The remaining answer books meant for evaluation shall be given only after ensuring that there is no significant variation in the marking of individual evaluators.
6	Evaluators will mark(\checkmark) wherever answer is correct. For wrong answer CROSS ‘X’ be marked. Evaluators will not put right (\checkmark)while evaluating which gives an impression that answer is correct and no marks are awarded. This is the most common mistake which evaluators are committing.
7	If a question has parts, please award marks on the right-hand side for each part. Marks awarded for different parts of the question should then be totaled up and written in the left-hand margin and encircled. This may be followed strictly.
8	If a question does not have any parts, marks must be awarded in the left-hand margin and encircled. This may also be followed strictly.
9	If a student has attempted an extra question, answer of the question deserving more marks should be retained and the other answer scored out with a note “Extra Question” . However, for MCQs(Q1 – Q20), only first attempt to be evaluated.
10	No marks to be deducted for the cumulative effect of an error. It should be penalized only once.

11	A full scale of marks _____(example 0 to 80/70/60/50/40/30 marks as given in Question Paper) has to be used. Please do not hesitate to award full marks if the answer deserves it.
12	Every examiner has to necessarily do evaluation work for full working hours i.e., 8 hours every day and evaluate 20 answer books per day in main subjects and 25 answer books per day in other subjects (Details are given in Spot Guidelines).
13	Ensure that you do not make the following common types of errors committed by the Examiner in the past:- <ul style="list-style-type: none"> ● Leaving answer or part thereof unassessed in an answer book. ● Giving more marks for an answer than assigned to it. ● Wrong totaling of marks awarded on an answer. ● Wrong transfer of marks from the inside pages of the answer book to the title page. ● Wrong question wise totaling on the title page. ● Wrong totaling of marks of the two columns on the title page. ● Wrong grand total. ● Marks in words and figures not tallying/not same. ● Wrong transfer of marks from the answer book to online award list. ● Answers marked as correct, but marks not awarded. (Ensure that the right tick mark is correctly and clearly indicated. It should merely be a line. Same is with the X for incorrect answer.) ● Half or a part of answer marked correct and the rest as wrong, but no marks awarded.
14	While evaluating the answer books if the answer is found to be totally incorrect, it should be marked as cross (X) and awarded zero (0)Marks.
15	Any un assessed portion, non-carrying over of marks to the title page, or totaling error detected by the candidate shall damage the prestige of all the personnel engaged in the evaluation work as also of the Board. Hence, in order to uphold the prestige of all concerned, it is again reiterated that the instructions be followed meticulously and judiciously.
16	The Examiners should acquaint themselves with the guidelines given in the “ Guidelines for spot Evaluation ” before starting the actual evaluation.
17	Every Examiner shall also ensure that all the answers are evaluated, marks carried over to the title page, correctly totaled and written in figures and words.
18	The candidates are entitled to obtain photocopy of the Answer Book on request on payment of the prescribed processing fee. All Examiners/Additional Head Examiners/Head Examiners are once again reminded that they must ensure that evaluation is carried out strictly as per value points for each answer as given in the Marking Scheme.


MARKING SCHEME (430/5/3)
MATHEMATICS (BASIC)

1. The sum of the first 50 odd natural numbers is :
 (a) 5000 (b) 2500
 (c) 2550 (d) 5050

Answer : (b) 2500

1

2. In the given figure, $AD = 2$ cm, $DB = 3$ cm, $DE = 2.5$ cm and $DE \parallel BC$. The value of x is:
 (a) 6cm (b) 3.75 cm
 (c) 6.25 cm (d) 7.5 cm



Answer : (c) 6.25 cm

1

3. A circle is of radius 3 cm. The distance between two of its parallel tangents is:
 (a) 12 cm (b) 6 cm
 (c) 3 cm (d) 4.5 cm

Answer: (b) 6 cm

1

4. The median class for the data given below is :

Class	20 – 40	40 – 60	60 – 80	80 – 100	100 – 120
Frequency	10	12	14	13	17

- (a) 80 – 100 (b) 20 – 40
 (c) 40 – 60 (d) 60 – 80

Answer: (d) 60 – 80

1

5. Mean and median of some data are 32 and 30 respectively. Using empirical relation, mode of the data is :

- (a) 36 (b) 26
 (c) 30 (d) 20

Answer : (b) 26

1

6. In two triangles ΔPQR and ΔABC , it is given that $\frac{AB}{BC} = \frac{PQ}{PR}$. For these two triangles to be similar, which of the following should be true ?

- (a) $\angle A = \angle P$ (b) $\angle B = \angle Q$
 (c) $\angle B = \angle P$ (d) $CA = QR$

Answer: (c) $\angle B = \angle P$

1

7. If $\sin \theta = \frac{3}{4}$, then $\frac{(\sec^2 \theta - 1) \cos^2 \theta}{\sin \theta}$ equals :

- (a) $\frac{3}{4}$ (b) $\frac{3}{4}$
 (c) $\frac{5}{4}$ (d) $\frac{9}{16}$
 (e) $\frac{3}{4}$


Answer : (b) $\frac{3}{4}$ 1

8. The 8th term of an A.P. is 17 and its 14th term is 29. The common difference of this A.P. is :

- (a) 3 (b) 2
 (c) 5 (d) -2

Answer : (b) 2 1

9. In the given figure, O is the centre of the circle and PA is a tangent to the circle. If $\angle OAB = 60^\circ$, then $\angle OPA$ is equal to :



- (a) 60° (b) 30°
 (c) 15° (d) 20°

Answer : (b) 30° 1

10. One card is drawn at random from a well shuffled deck of 52 playing cards. The probability that it is a red king is :

- (a) $\frac{1}{52}$ (b) $\frac{1}{26}$
 (c) $\frac{2}{26}$ (d) $\frac{2}{13}$

Answer: (b) $\frac{1}{26}$ 1

11. If the lines represented by equations $3x + 2my = 2$ and $2x + 5y + 1 = 0$ are parallel, then the value of m is :

- (a) $\frac{2}{5}$ (b) $-\frac{5}{4}$
 (c) $\frac{3}{2}$ (d) $\frac{15}{4}$

Answer: (d) $\frac{15}{4}$

1

12. $\Delta ABC \sim \Delta DEF$ and their perimeter are 32 cm and 24 cm respectively.

If $AB = 10$ cm, then DE equals :

- (a) 8 cm (d) 7.5 cm
(c) 15 cm (d) $5\sqrt{3}$ cm

Answer: (b) 7.5 cm

1

13. The two roots of the equation $3x^2 - 2\sqrt{6}x + 2 = 0$ are:

- (a) real and distinct
(b) not real
(c) real and equal
(d) rational

Answer: (c) real and equal

1

14. If $\sin \theta = \frac{a}{b}$, then $\sec \theta$ is equal to ($0 \leq \theta \leq 90^\circ$):

- (a) $\frac{a}{\sqrt{b^2 - a^2}}$ (b) $\frac{b}{\sqrt{b^2 - a^2}}$
(c) $\frac{\sqrt{b^2 - a^2}}{b}$ (d) $\frac{\sqrt{b^2 - a^2}}{a}$

Answer: (b) $\frac{b}{\sqrt{b^2 - a^2}}$

1

15. The distance between the points A (0, 6) and B (-6, 2) is :

- (a) 6 units (b) $2\sqrt{6}$ units
(c) $2\sqrt{13}$ units (d) $13\sqrt{2}$ units

Answer : (c) $2\sqrt{13}$ units

1

16. The value(s) of k for which the roots of quadratic equation $x^2 + 4x + k = 0$ are real, is :

- (a) $k \geq 4$ (b) $k \leq 4$
(c) $k \geq -4$ (d) $k \leq -4$

Answer: (b) $k \leq 4$

1

17. HCF of $(3^4 \times 2^2 \times 7^3)$ and $(3^2 \times 5 \times 7)$ is :

- (a) 630 (b) 63
(c) 729 (d) 567

Answer: (b) 63

1

18. If one zero of the quadratic polynomial $kx^2 + 3x + k$ is 2, then the value of k is:

- (a) $-\frac{6}{5}$ (b) $\frac{6}{5}$
(b) $\frac{5}{6}$ (d) $-\frac{5}{6}$

Answer: (a) $-\frac{6}{5}$

1

Questions number 19 and 20 are Assertion and Reason based questions carrying 1 mark each. Two statements are given, one labelled as Assertion (A) and the other is labelled as Reason (R). Select the correct answer to these questions from the codes (a), (b), (c) and (d) as given below.

- (a) Both Assertion (A) and Reason (R) are true and Reason (R) is the correct explanation of the Assertion (A).
- (b) Both Assertion (A) and Reason (R) are true, but Reason (R) is **not** the correct explanation of the Assertion (A).
- (c) Assertion (A) is true, but Reason (R) is false.
- (d) Assertion (A) is false, but Reason (R) is true.

19. **Assertion (A)** : The surface area of largest sphere that can be inscribed in a hollow cube of side 'a' cm is $\pi a^2 \text{ cm}^2$.

Reason (R) : The surface area of a sphere of radius 'r' is $\frac{4}{3}\pi r^3$.

Answer: (c) Assertion (A) is true, but Reason (R) is false.

1

20. **Assertion (A)** : When two coins are tossed together, the probability of getting no tail is $\frac{1}{4}$.


Reason (R) : The probability P(E) of an event E satisfies $0 \leq P(E) \leq 1$.

Answer: (b) Both Assertion (A) and Reason (R) are true, but Reason (R) is **not** the correct explanation of the Assertion (A).

1

SECTION B

21. In the given figure, tangents AB and AC are drawn to a circle centred at O. If $\angle OAB = 60^\circ$ and $OB = 5 \text{ cm}$, find lengths OA and AC.



Solution. $\sin 60^\circ = \frac{\sqrt{3}}{2} = \frac{5}{OA} \Rightarrow OA = \frac{10\sqrt{3}}{3} \text{ cm}$

$$\frac{1}{2} + \frac{1}{2}$$

$$\tan 60^\circ = \sqrt{3} = \frac{5}{AB} \Rightarrow AB = \frac{5\sqrt{3}}{3} \text{ cm} = AC$$

$$\frac{1}{2} + \frac{1}{2}$$

22. Find the sum of the first 20 terms of the A.P. : $-\frac{29}{3}, -9, -\frac{25}{3}, -\frac{23}{3}, \dots$

Solution: $d = \frac{2}{3}$

$$\frac{1}{2}$$

$$\therefore S_{20} = \frac{20}{2} \left[-\frac{58}{3} + 19 \times \frac{2}{3} \right]$$

$$1$$

$$= -\frac{200}{3}$$

$$\frac{1}{2}$$

23. (a) Evaluate: $\frac{\sin 30^\circ + \tan 45^\circ}{\sec 30^\circ + \cot 45^\circ}$

Solution: (a) Required value is $\frac{\frac{1}{2} + 1}{\frac{2}{\sqrt{3}} + 1} = \frac{3\sqrt{3}}{2(2 + \sqrt{3})}$ $1 \frac{1}{2}$
 $\frac{1}{2}$

OR

(b) For $A = 30^\circ$ and $B = 60^\circ$, verify that :

$$\sin(A + B) = \sin A \cos B + \cos A \sin B.$$

Solution (b) LHS = $\sin 90^\circ = 1$ $\frac{1}{2}$

RHS = $\frac{1}{2} \times \frac{1}{2} + \frac{\sqrt{3}}{2} \times \frac{\sqrt{3}}{2}$ 1

= 1 $\frac{1}{2}$

LHS = RHS

24. Find LCM of 480 and 256 using prime factorization.

Solution: $480 = 2^5 \times 3 \times 5$ $\frac{1}{2}$

$256 = 2^8$ $\frac{1}{2}$

LCM (480, 256) = $2^8 \times 3 \times 5 = 3840$ 1

25. (a) Show that A(1, 2), B(5, 4), C(3, 8) and D(-1, 6) are vertices of a parallelogram ABCD.

Solution: Mid-point of AC = $\left(\frac{3+1}{2}, \frac{8+2}{2}\right) = (2, 5)$ 1

Mid-point of BD = $\left(\frac{5-1}{2}, \frac{4+6}{2}\right) = (2, 5)$ $\frac{1}{2}$

\therefore Mid-point of AC = Mid-point of BD $\frac{1}{2}$

Hence, ABCD is a parallelogram

OR

(b) Show that the points A(3, 0), B(6, 4) and C(-1, 3) are the vertices of a right angled triangle.

Solution: $AB^2 = 3^2 + 4^2 = 25$ $\frac{1}{2}$

$BC^2 = 7^2 + 1^2 = 50$ $\frac{1}{2}$

$$AC^2 = 4^2 + 3^2 = 25$$

$\frac{1}{2}$
 $\frac{1}{2}$

$$\therefore BC^2 = AB^2 + AC^2$$

$\therefore \Delta ABC$ is right-angled triangle.

SECTION C

26. Find mean of the following data :

Class	0 – 15	15 – 30	30 – 45	45 – 60	60 – 75	75 – 90
Frequency	12	15	11	20	16	6

Solution.

Class	x	f	$u = \frac{x - 37.5}{15}$	fu
0 – 15	7.5	12	-2	-24
15 – 30	22.5	15	-1	-15
30 – 45	37.5	11	0	0
45 – 60	52.5	20	1	20
60 – 75	67.5	16	2	32
75 – 90	82.5	6	3	18
		80		31


For Correct Table: 2 Marks

$$\begin{aligned} \text{Mean} &= a + \frac{\sum fu}{\sum f} \times h \\ &= 37.5 + 15 \times \frac{31}{80} = 43.3 \end{aligned}$$

1

27. (a) Determine the ratio in which the point P(a, -2) divides the line segment joining the points A(-4, 3) and B(2, -4). Also, find the value of 'a'.

Solution:



$$\text{Let } AP : PB = k : 1$$

$$\therefore \frac{-4k + 3}{k + 1} = -2 \Rightarrow k = \frac{5}{2}$$


$$\text{So, } AP : PB = 5 : 2$$

$$\text{Hence, } \frac{10 - 8}{7} = a \Rightarrow a = \frac{2}{7}$$

$1 + \frac{1}{2}$
 $\frac{1}{2}$
1

OR

- (b) In the given figure, in ΔABC points D and E are mid-points of sides BC and AC respectively. If given vertices are $A(4, -2)$, $B(2, -2)$ and $C(-6, -7)$, then verify the result $DE = \frac{1}{2} AB$.



Solution: Point D is $\left(-2, \frac{-9}{2}\right)$ 1

Point E is $\left(-1, \frac{-9}{2}\right)$ 1

$\therefore DE = \sqrt{1^2 + 0^2} = 1$ and $AB = \sqrt{2^2 + 0^2} = 2$ $\frac{1}{2} + \frac{1}{2}$

$\therefore DE = \frac{1}{2} AB$

28. ABC is an isosceles triangle with $AB = AC$, circumscribed about a circle. Prove that BC is bisected at E.


Solution. $AD = AF$, $BD = BE$ and $CE = CF$ (tangents from external point) 1

$AB = AC \Rightarrow AD + DB = AF + FC$ $\frac{1}{2}$

$\Rightarrow AF + DB = AF + FC$ $\frac{1}{2}$

$\Rightarrow DB = FC$ $\frac{1}{2}$

$\Rightarrow BE = EC$ or BC is bisected at E. $\frac{1}{2}$



29. Prove that : $\sqrt{\frac{\sec\theta-1}{\sec\theta+1}} + \sqrt{\frac{\sec\theta+1}{\sec\theta-1}} = 2\operatorname{cosec} \theta$

Solution: LHS = $\frac{\sqrt{\sec\theta-1}}{\sqrt{\sec\theta+1}} + \frac{\sqrt{\sec\theta+1}}{\sqrt{\sec\theta-1}} = \frac{2 \sec \theta}{\sqrt{\sec^2 \theta - 1}}$ 1

= $\frac{2 \sec \theta}{\tan \theta}$ 1

= $\frac{2 \cos \theta}{\cos \theta \cdot \sin \theta} = 2 \operatorname{cosec} \theta = \text{RHS}$ 1

30. (a) Sabina went to a bank ATM to withdraw ₹2,000. She received ₹ 50 and ₹ 100 notes only. If Sabina got 25 notes in all, how many notes of ₹ 50 and ₹ 100 did she receive?

Solution: (a) Let number of ₹50 notes be x and number of ₹100 notes be y .

ATQ, $x + y = 25$ _____ (i) 1

And, $50x + 100y = 2000$ _____ (ii) 1

Solving (i) and (ii), $x = 10$, $y = 15$ 1

Number of ₹50 notes = 10 and Number of ₹100 notes = 15

OR

- (b) Five years ago, Amit was thrice as old as Baljeet. Ten years hence, Amit shall be twice as old as Baljeet. What are their present ages?

Solution : (b) Let Amit's present age be x years and Baljeet's present age be y years.

ATQ, $(x - 5) = 3(y - 5) \Rightarrow x - 3y = -10$ 1

And $(x + 10) = 2(y + 10) \Rightarrow x - 2y = 10$ 1

Solving equations to get, $y = 20$, $x = 50$ 1

Amit's present age = 50 years and Baljeet's present age = 20 years

31. Prove that $11 + 3\sqrt{2}$ is an irrational number, given that $\sqrt{2}$ is an irrational number.

Solution: Let us assume that $11 + 3\sqrt{2}$ be a rational number.

$\Rightarrow 11 + 3\sqrt{2} = \frac{a}{b}$, where a and b are integers, $b \neq 0$ 1

$\Rightarrow \sqrt{2} = \frac{a - 11b}{3b}$ 1

RHS is a rational number but LHS is irrational. 1

\therefore Our assumption was wrong. Hence, $11 + 3\sqrt{2}$ is irrational. } 1

SECTION D

32. (a) Divide 16 into two parts such that twice the square of the greater part, exceeds the square of the smaller part by 164

Solution: (a) Let the greater part be x then the other part is $16 - x$

ATQ, $2x^2 = (16 - x)^2 + 164$ 1

$\Rightarrow x^2 + 32x - 420 = 0$ 1

$\Rightarrow (x + 42)(x - 10) = 0$ 1

$\Rightarrow x = -42$, $x = 10$ 1

$\Rightarrow x = 10$ ($x \neq -42$) 1

\therefore Two parts are 10 and 6. 1

OR

- (b) A motor boat whose speed is 18 km/h in still water takes 1 hour more to go 24 km upstream, than to return to the same point. Find the speed of the stream and total time of the journey.

Solution: Let the speed of the stream be x km/hour.

ATQ, $\frac{24}{18 - x} - \frac{24}{18 + x} = 1$ 2

$$\Rightarrow x^2 + 48x - 324 = 0 \quad 1$$


$$\Rightarrow (x + 54)(x - 6) = 0$$

$$\Rightarrow x = 6 \quad (x \neq -54) \quad 1$$

$$\text{Total time of journey} = \frac{24}{12} + \frac{24}{24} = 3 \text{ hours} \quad 1$$

33. If a line is drawn parallel to one side of a triangle to intersect the other two sides at distinct points, then prove that the other two sides are divided in the same ratio.

Solution.



For figure:
1 mark

Given: In $\triangle ABC$, $DE \parallel BC$

$\frac{1}{2}$

To prove: $\frac{AD}{DB} = \frac{AE}{EC}$

$\frac{1}{2}$

Const.: Join BE , CD . Draw $DM \perp AC$, $EN \perp AB$

$\frac{1}{2}$

Proof: $\frac{\text{ar}(\triangle ADE)}{\text{ar}(\triangle BDE)} = \frac{\frac{1}{2} \times AD \times EN}{\frac{1}{2} \times DB \times EN} = \frac{AD}{DB}$ _____ (i)

1

Similarly, $\frac{\text{ar}(\triangle ADE)}{\text{ar}(\triangle CDE)} = \frac{AE}{EC}$ _____ (ii)

$\frac{1}{2}$

$\triangle BDE$ and $\triangle CDE$ are on the same base DE and between the same parallel lines BC and DE .

$\text{ar}(\triangle BDE) = \text{ar}(\triangle CDE)$ _____ (iii)


$\frac{1}{2}$

From (i), (ii) and (iii) $\frac{AD}{DB} = \frac{AE}{EC}$

$\frac{1}{2}$

34. From the top of a building 50 m high, the angles of depression of the top and bottom of a tower are observed to be 30° and 60° . Find the height of the tower and distance between the building and the tower. (Take $\sqrt{3} = 1.73$)

Solution:



For Figure:
1 mark

Let AB be the building and CD be the tower.


$$\text{In } \triangle BED, \quad \tan 30^\circ = \frac{50-h}{x} \Rightarrow 50-h = \frac{x}{\sqrt{3}} \quad \text{----- (i)} \quad 1 + \frac{1}{2}$$

$$\text{In } \triangle BAC, \quad \tan 60^\circ = \frac{50}{x} \Rightarrow x = \frac{50}{\sqrt{3}} \quad \text{----- (ii)} \quad 1 + \frac{1}{2}$$

$$\text{Solving (i) and (ii) to get } h = \frac{100}{3} \text{ or } 33.33 \text{ m} \quad \frac{1}{2}$$

$$\text{and } x = 28.83 \text{ m} \quad \frac{1}{2}$$

35. (a) In the given figure, AB is a chord of a circle of radius 7 cm and centred at O. Find the area of the shaded region if $\angle AOB = 90^\circ$. Also, find length of minor arc AB.



$$\begin{aligned} \text{Solution: Area of sector AOB} &= \frac{22}{7} \times 7 \times 7 \times \frac{90}{360} && 1 \\ &= \frac{77}{2} \text{ cm}^2 && \frac{1}{2} \end{aligned}$$


$$\text{Area of } \triangle AOB = \frac{1}{2} \times 7 \times 7 = \frac{49}{2} \text{ cm}^2 \quad 1$$

$$\therefore \text{ Shaded area} = \frac{77}{2} - \frac{49}{2} = \frac{28}{2} = 14 \text{ cm}^2 \quad 1$$

$$\text{Length of arc AB} = 2 \times \frac{22}{7} \times 7 \times \frac{90}{360} = 11 \text{ cm} \quad 1 + \frac{1}{2}$$

OR

- (b) AB and CD are arcs of two concentric circles of radii 3.5 cm and 10.5 cm respectively and centered at O. Find the area of the shaded region if $\angle AOB = 60^\circ$. Also, find the length of arc CD.



Solution: (b) Here OA = 3.5 cm, OC = 10.5 cm

$$\begin{aligned} \text{Shaded area} &= \pi \times \frac{60}{360} (10.5^2 - 3.5^2) && 2 \\ &= \frac{22}{7} \times \frac{1}{6} \times 98 && 1 \end{aligned}$$

$$= \frac{154}{3} \text{ cm}^2 \text{ or } 51.3 \text{ cm}^2 \quad \frac{1}{2}$$

$$\text{Length of arc CD} = 2 \times \frac{22}{7} \times 10.5 \times \frac{60}{360} \quad 1$$

$$= 11 \text{ cm} \quad \frac{1}{2}$$

SECTION E

36. Singing bowls (hemispherical in shape) are commonly used in sound healing practices. Mallet (cylindrical in shape) is used to strike the bowl in a sequence to produce sound and vibration.



One such bowl is shown here whose dimensions are :

Hemispherical bowl has outer radius 6 cm and inner radius 5 cm.

Mallet has height of 10 cm and radius 2 cm.

Based on the above, answer the following questions :


- (i) What is the volume of the material used in making the mallet ?
- (ii) The bowl is to be polished from inside. Find the inner surface area of the bowl.
- (iii) (a) Find the volume of metal used to make the bowl.

OR

- (iii) (b) Find total surface area of the mallet. (Use $\pi = 3.14$)

Solution:	(i) Volume of material = $3.14 \times 2 \times 2 \times 10 = 125.6 \text{ cm}^3$	1
	(ii) Inner SA of the bowl = $2 \times 3.14 \times 25 = 157 \text{ cm}^2$	1
	(iii) (a) Volume of the metal = $\frac{2}{3} \times 3.14 \times (6^3 - 5^3)$	1
	= 190.5 cm^3	1
	OR	
	(iii) (b) Total SA of mallet = $2 \times 3.14 \times 2 (2 + 10)$	1
	= 150.7 cm^2	1

37. Rainbow is an arch of colours that is visible in the sky after rain or when water droplets are present in the atmosphere. The colours of the rainbow are generally, red, orange, yellow, green, blue, indigo and violet. Each colour of the rainbow makes a parabola. We know that any quadratic polynomial $p(x) = ax^2 + bx + c$ ($a \neq 0$) represents a parabola on the graph paper.



Based on the above, answer the following questions :

- (i) The graph of a rainbow $y = f(x)$ is shown in the figure. Write the number of zeroes of the curve.
- (ii) If the graph of a rainbow does not intersect the x-axis but intersects y-axis at one point, then how many zeroes will it have ?
- (iii) (a) If a rainbow is represented by the quadratic polynomial $p(x) = x^2 + (a + 1)x + b$, whose zeroes are 2 and -3 , find the value of a and b.

OR

- (iii) (b) The polynomial $x^2 - 2x - (7p + 3)$ represents a rainbow. If -4 is a zero of it, find the value of p.


Solution.

	(i) Two zeroes	1
	(ii) 0 or no zero	1
	(iii) (a) Getting $2a+b = -6$ and $-3a+b = -6$	$\frac{1}{2} + \frac{1}{2}$
	Solving to get $a = 0$ and $b = -6$	$\frac{1}{2} + \frac{1}{2}$

OR

- | | | |
|--|--|---|
| | (iii) (b) -4 is a zero of the given polynomial $\Rightarrow 21-7p = 0$ | 1 |
| | $\Rightarrow p = 3$ | 1 |
-

38. Some students were asked to list their favourite colour. The measure of each colour is shown by the central angle of a pie chart given below :



Study the pie chart and answer the following questions :

- (i) If a student is chosen at random, then find the probability of his/her favourite colour being white ?
- (ii) What is the probability of his/her favourite colour being blue or green ?
- (iii) (a) If 15 students liked the colour yellow, how many students participated in the survey ?

OR

- (iii) (b) What is the probability of the favourite colour being red or blue ?

Solution. (i) $P(\text{favourite colour being white}) = \frac{120}{360}$ or $\frac{1}{3}$ 1

(ii) $P(\text{favourite colour being blue or green}) = \frac{60 + 60}{360}$ or $\frac{1}{3}$ 1

(iii) (a) Let total number of students be x

$$\Rightarrow \frac{15}{x} = \frac{1}{4} \quad \text{1} \frac{1}{2}$$

$$\Rightarrow x = 60 \quad \frac{1}{2}$$

OR

(iii)(b) $P(\text{favourite colour being red or blue}) = \frac{60 + 30}{360} = \frac{1}{4}$ $1+1$